

PULMONARY FUNCTION ELIGIBILITY FORM

ID FORM CODE: PFT Visit SEQ VERSION: 1.0 10/26/10 Number #
0a) Form Date
<u>Instructions:</u> This form should be completed prior to conducting pulmonary function testing. Whenever numerical responses are required, enter the number so that the last digit appears in the rightmost box. Enter leading zeroes where necessary to fill all boxes.
1) Have you eaten a large meal with the last 2 hours?
Yes
NON
2) Have you smoked within the last hour?
Yes
3) Have you participated in any vigorous exercise in the past 30 minutes?
No
4) Have you consumed any alcoholic beverages within the past 4 hours?
Yes
NO
5) Have you taken any inhaled or oral medication for your lungs in the past 48 hours?
Yes
6) Have you used tiotropium (Spiriva) within in the past 48 hours?
No
6a) Date//
7) Have you used theophylline (Theo 24, Uniphyl, Theochron) within past 48 hrs?
Yes Y
100
No

ID NUMBER: FORM CODE: PFT Visit VERSION: 1.0 10/26/10 Number SEQ #									
2) an extended release preparation (Theochron, generic SR)3) an immediate release preparation (Elixophyllin, Theolair)									
7b) Date// time: AM/PM (circle one)									
8) Have you used another once-a-day bronchodilator within in the past 48 hours?									
No									
8a) Date/ time: AM/PM (circle one) 8b) Name:									
9) Have you used any long-acting (12-hour) beta agonist within in the past 24 hours?									
YesY									
NoN →Go to item10									
9a) What is the most recent long acting beta agonist you have used?:									
1) Albuterol sustained release tab. (VoSpire ER):									
2) Arformoterol (Brovana):									
3) Formoterol (Dulera, Foradil, Perforomist, Symbicort):									
4) Salmeterol (Advair Diskus, Advair HFA, Serevent Diskus) 5) Other 12 hour beta agonist: Name:									
9b) Time AM/PM (circle one)									
10) Have you used ipratropium within the past 8 hours?									
YesY									
No									
10a) What is the most recent ipratropium or ipratropium/albuterol combination have you used?									
I) Ipratropium (Atrovent, generic ipratropium nebulizer solution)									
2) Ipratropium/albuterol combination (Combivent, Duoneb, generic nebulizer solution)									
10b) Time AM/PM (circle one)									
If <300 minutes since last ipratropium dose, see partial dosing instruction in the PFT MOP.									

ID NUMBER:								FORM CODE: PFT VERSION: 1.0 10/26/10	Visit Number		SEQ#	
l1) Have you υ	ısed	any s	shor	t-acti	ng be	ta ago	onist	within in the past 6 ho	ours?]
Yes		Y										
No												
11a) What	is the	e mo:	st re	cent	short	-actin	g beta	a agonist have you us	ed?			
1) Alb releas					, Prov	entil l	HFA,	Ventolin HFA, Acune	b, generic	solutio	n, non-susta	ined
2) Epi	neph	rine	(Epi	pen,	Twinj	ect, S	2, ge	neric)				
3) Lev	albut	terol	(Xor	oene:	x HF	A, Xop	enex	solution, generic solu	ution)			
4) Metaproterenol (generic solution, syrup, non-sustained release tablets)												
5) Pirbuterol (Maxair)												
6) Terbutaline (generic tablets)												
			_				Jame	9				
7) 011					a age	// IIOt. 1	varric	/				
11b) Time		_]:[AM/P	M (ciro	cle oi	ne)				
If <165 minut	es s	ince	last	sho	rt-act	ing b	eta a	gonist dose, see pa	rtial dosii	ng instr	ructions in	PFT MOP.
l2) Have you h	nad m	nore	then	18 c	z. of	coffee	or e	equivalent caffeine, 20	00 mg), in	the pas	st 6 hours?	
Yes							Υ					
No							N					

200 mg= (roughly):18 oz of coffee, 4 shots of expresso, 40 oz of tea, 60 oz of a cola soft drink, 16 oz of energy drink